

On some classical and bipartite Ramsey numbers

Janusz Dybizbański, Tomasz Dzido

Institute of Informatics, University of Gdańsk

Wita Stwosza 57, 80-952 Gdańsk, Poland

Abstract

In talk we present definitions of classical and bipartite Ramsey numbers. We show the results for example $R(C_4, C_4, K_4 - e) = 16$, which fill few of the gaps in the tables presented in [1]. Moreover by using computer methods we improve lower and upper bounds for some another multicolor classical and bipartite Ramsey numbers involving quadrilateral C_4 and other graphs. Our results improve bounds presented in [4], [5] and [2].

References

- [1] Arste J., Klamroth K., Mengersen I.: Three color Ramsey numbers for small graphs, *Utilitas Mathematica* **49** (1996) 85–96.
- [2] Goddard W., Henning M.A., Oellermann O.R.: Bipartite Ramsey numbers and Zarankiewicz numbers, *Discrete Mathematics* **219** (2000)
- [3] Radziszowski S.P.: Small Ramsey Numbers, *Electronic Journal of Combinatorics*, Dynamic Survey 1, revision #12, August 2009, <http://www.combinatorics.org>.
- [4] Radziszowski S.P., Shao Z., Xu X.: Bounds on some Ramsey numbers involving quadrilateral, *Ars Combinatoria* **90** (2009) 337–344.
- [5] Radziszowski S.P., Xu X.: $28 \leq R(C_4, C_4, C_3, C_3) \leq 36$, *Utilitas Mathematica* **79** (2009) 353–357.